



Active Bacterial Core Surveillance (ABCs) Report

Emerging Infections Program Network

group B streptococcus, 2003 – *preliminary*



ABCs Areas

California (3 county San Francisco Bay area); Colorado (5 county Denver area); Connecticut; Georgia (20 county Atlanta area); Maryland; Minnesota; New York (7 county Rochester area and 8 county Albany area); Oregon (3 county Portland area); Tennessee (11 urban counties)

ABCs Population

The surveillance areas represent 30,242,564 persons and 425,405 persons <1 yr. Source: National Center for Health Statistics bridged-race vintage 2002 postcensal file

ABCs Case Definition

Invasive group B streptococcal disease: isolation of group B streptococcus from a normally sterile site in a resident of a surveillance area in 2003. Early-onset cases occur at <7 days of age and late-onset occur between 7 and 90 days of age.

ABCs Methodology

ABCs personnel routinely contacted all microbiology laboratories serving acute care hospitals in their area to identify cases. Standardized case report forms that include information on demographic characteristics, clinical syndrome, and outcome of illness were completed for each identified case. Regular laboratory audits assessed completeness of active surveillance and detected additional cases.

Rates of invasive group B streptococcal disease were calculated using population estimates for 2002. Rates of early-onset and late-onset group B streptococcal disease were calculated using population estimates for persons <1 yr for 2002. For national projections of cases, race- and age-specific rates of disease were applied from the aggregate surveillance area to the age and racial distribution of the 2002 U. S. population. Cases with unknown race were distributed by area based on reported race distribution for known cases within the eight age categories.

Reported ABCs Profiles

Race	No. (Rate*)	
White	1,468	(6.3)
Black	564	(11.3)
Other	57	(2.9)
Unknown race (n=257) distributed among knowns.		
*Cases per 100,000 population for ABCs areas.		
Ethnicity	No. (Rate*)	
Hispanic	108	(4.4)
Non-Hispanic	901	-----
Unknown	1,071	-----

*Cases per 100,000 population for ABCs areas.

Age (years)	Cases	Deaths
	No. (Rate*)	No. (Rate*)
<1	287(67.5)	12(2.8)
1	2(0.5)	0(0.0)
2-4	0(0.0)	0(0.0)
5-17	13(0.2)	0(0.0)
18-34	140(1.9)	7(0.1)
35-49	308(4.2)	23(0.3)
50-64	491(10.3)	53(1.1)
≥65	848(25.4)	123(3.7)
Total	2,089(6.9)	218(0.7)

*Cases or deaths per 100,000 population for ABCs areas

Race	Early-Onset	Late-Onset
	No. (Rate*)	No. (Rate*)
White	81(0.27)	71(0.23)
Black	44(0.53)	58(0.71)
Other	5(0.20)	3(0.10)
Total	130(0.31)	132(0.31)

Unknown race (n=19 Early-Onset, n=14 Late-Onset) distributed among knowns.

*Cases per 1,000 live births for ABCs areas

National Projection of Invasive Disease

Early-Onset Cases: 1,250 (0.31/1,000 live births)
Late-Onset Cases: 1,250 (0.31/1,000 live births)
Total Cases: 20,150 (7.0/100,000 population)
Deaths: 2,100 (0.7/100,000 population)

Healthy People 2010 Update

Early-Onset Disease

Objective: Decrease the incidence of invasive early-onset group B streptococcal disease to 0.5 cases per 1,000 live births.

Race	2010 Objective	2003 Rate*
White	0.5/1,000	0.27/1,000
Black	0.5/1,000	0.53/1,000
Other	0.5/1,000	0.20/1,000
Total	0.5/1,000	0.31/1,000

*Projected cases per 1,000 live births

For more information, visit our web sites

<http://www.cdc.gov/ncidod/dbmd/abcs>

<http://www.cdc.gov/groupbstrep>

Centers for Disease Control and Prevention. 2004. Active Bacterial Core Surveillance Report, Emerging Infections Program Network, Group B Streptococcus, 2003 – *provisional*. Available via the Internet:

<http://www.cdc.gov/ncidod/dbmd/abcs/survreports/gbs02.pdf>